

On Hermite-Hadamard Inequality for Twice Differentiable Functions Bounded by Exponentials

S. S. Dragomir^{[a],[b],*} and I. Gomm^[a]

^[a] Mathematics, School of Engineering & Science, Victoria University, Melbourne City, MC 8001, Australia.

^[b] School of Computational & Applied Mathematics, University of the Witwatersrand, Private Bag 3, Johannesburg 2050, South Africa.

* Corresponding author.

Address: Mathematics, School of Engineering & Science, Victoria University, PO Box 14428, Melbourne City, 8001, Australia; E-Mail: sever.dragomir@vu.edu.au

Received: September 17, 2012/ Accepted: November 2, 2012/ Published: November 30, 2012

Abstract: Some Hermite-Hadamard type inequalities for twice differentiable functions whose second derivatives are bounded below and above by exponentials are given. Applications for special means are provided as well.

Key words: Convex functions; Hermite-Hadamard inequality; Special means

Dragomir, S. S., & Gomm, I. (2012). On Hermite-Hadamard Inequality for Twice Differentiable Functions Bounded by Exponentials. *Studies in Mathematical Sciences*, 5(2), 44–54. Available from <http://www.cscanada.net/index.php/sms/article/view/j.sms.1923845220120502.2583> DOI: 10.3968/j.sms.1923845220120502.2583

1. INTRODUCTION

The following integral inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a)+f(b)}{2}, \quad (1.1)$$

which holds for any convex function $f : [a, b] \rightarrow \mathbb{R}$, is well known in the literature as the Hermite-Hadamard inequality.

There is an extensive amount of literature devoted to this simple and nice result which has many applications in the Theory of Special Means and in Information

Theory for divergence measures, from which we would like to refer the reader to the papers [1]-[58] and the references therein.

In this paper we establish some Hermite-Hadamard type inequalities for twice differentiable functions whose second derivatives are bounded below and above by exponential functions. Applications for special means are provided as well.

2. THE RESULTS

The following result holds:

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable function with the property that there exists the constants $\alpha, m, M \in \mathbb{R}$ with $\alpha \neq 0, m < M$ and such that*

$$me^{\alpha t} \leq f''(t) \leq Me^{\alpha t} \quad (2.1)$$

for any $t \in (a, b)$.

Then we have the inequalities

$$\begin{aligned} \frac{m}{\alpha^2} \left(\frac{e^{\alpha a} + e^{\alpha b}}{2} - \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} \right) \\ \leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \\ \leq \frac{M}{\alpha^2} \left(\frac{e^{\alpha a} + e^{\alpha b}}{2} - \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} \right) \end{aligned} \quad (2.2)$$

and

$$\begin{aligned} \frac{m}{\alpha^2} \left(\frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - e^{\alpha(\frac{a+b}{2})} \right) \\ \leq \frac{1}{b-a} \int_a^b f(t) dt - f\left(\frac{a+b}{2}\right) \\ \leq \frac{M}{\alpha^2} \left(\frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - e^{\alpha(\frac{a+b}{2})} \right). \end{aligned} \quad (2.3)$$

Proof. Consider the auxiliary function $g_{m,\alpha} : [a, b] \rightarrow \mathbb{R}$ given by $g_{m,\alpha}(t) := f(t) - \frac{m}{\alpha^2}e^{\alpha t}$. This function is twice differentiable and since $g_{m,\alpha}''(t) := f''(t) - me^{\alpha t} \geq 0$ we have that $g_{m,\alpha}$ is convex.

By the definition of convexity we have that

$$\begin{aligned} 0 &\leq \lambda g_{m,\alpha}(a) + (1-\lambda)g_{m,\alpha}(b) - g_{m,\alpha}(\lambda a + (1-\lambda)b) \\ &= \lambda f(a) + (1-\lambda)f(b) - f(\lambda a + (1-\lambda)b) \\ &\quad - \frac{m}{\alpha^2} \left(\lambda e^{\alpha a} + (1-\lambda)e^{\alpha b} - e^{\alpha(\lambda a + (1-\lambda)b)} \right) \end{aligned} \quad (2.4)$$

for any $\lambda \in [0, 1]$.

This is equivalent with

$$\begin{aligned} \frac{m}{\alpha^2} \left(\lambda e^{\alpha a} + (1-\lambda)e^{\alpha b} - e^{\alpha(\lambda a + (1-\lambda)b)} \right) \\ \leq \lambda f(a) + (1-\lambda)f(b) - f(\lambda a + (1-\lambda)b) \end{aligned} \quad (2.5)$$

for any $\lambda \in [0, 1]$.

Utilising the auxiliary function $g_{M,\alpha} : [a, b] \rightarrow \mathbb{R}$ given by $g_{M,\alpha}(t) := \frac{M}{\alpha^2} e^{\alpha t} - f(t)$ we also get

$$\begin{aligned} & \lambda f(a) + (1 - \lambda) f(b) - f(\lambda a + (1 - \lambda) b) \\ & \leq \frac{M}{\alpha^2} \left(\lambda e^{\alpha a} + (1 - \lambda) e^{\alpha b} - e^{\alpha(\lambda a + (1 - \lambda) b)} \right) \end{aligned} \quad (2.6)$$

for any $\lambda \in [0, 1]$.

Integrating the inequality (2.5) over $\lambda \in [0, 1]$ and taking into account that

$$\int_0^1 e^{\alpha(\lambda a + (1 - \lambda) b)} d\lambda = \frac{1}{b - a} \int_a^b e^{\alpha s} ds = \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b - a)}$$

and

$$\int_0^1 f(\lambda a + (1 - \lambda) b) d\lambda = \frac{1}{b - a} \int_a^b f(t) dt$$

we obtain the first inequality in (2.2).

The second part of (2.2) follows by (2.6) in the same way.

Now if we use (2.5) and (2.6) for $\lambda = \frac{1}{2}$ we get

$$\begin{aligned} \frac{m}{\alpha^2} \left(\frac{e^{\alpha u} + e^{\alpha v}}{2} - e^{\alpha(\frac{u+v}{2})} \right) & \leq \frac{f(u) + f(v)}{2} - f\left(\frac{u+v}{2}\right) \\ & \leq \frac{M}{\alpha^2} \left(\frac{e^{\alpha u} + e^{\alpha v}}{2} - e^{\alpha(\frac{u+v}{2})} \right) \end{aligned} \quad (2.7)$$

for any $u, v \in [a, b]$.

If we write this inequality for $u = \lambda a + (1 - \lambda) b$ and $v = (1 - \lambda) a + \lambda b$ then we get

$$\begin{aligned} & \frac{m}{\alpha^2} \left(\frac{e^{\alpha(\lambda a + (1 - \lambda) b)} + e^{\alpha((1 - \lambda) a + \lambda b)}}{2} - e^{\alpha(\frac{a+b}{2})} \right) \\ & \leq \frac{f(\lambda a + (1 - \lambda) b) + f((1 - \lambda) a + \lambda b)}{2} - f\left(\frac{a+b}{2}\right) \\ & \leq \frac{M}{\alpha^2} \left(\frac{e^{\alpha(\lambda a + (1 - \lambda) b)} + e^{\alpha((1 - \lambda) a + \lambda b)}}{2} - e^{\alpha(\frac{a+b}{2})} \right) \end{aligned} \quad (2.8)$$

for any $\lambda \in [0, 1]$.

Integrating the inequality (2.8) over λ on the interval $[0, 1]$ and taking into account that

$$\int_0^1 e^{\alpha(\lambda a + (1 - \lambda) b)} d\lambda = \int_0^1 e^{\alpha((1 - \lambda) a + \lambda b)} d\lambda = \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b - a)}$$

and

$$\begin{aligned} \int_0^1 f(\lambda a + (1 - \lambda) b) d\lambda & = \int_0^1 f((1 - \lambda) a + \lambda b) d\lambda \\ & = \frac{1}{b - a} \int_a^b f(t) dt \end{aligned}$$

then we get the desired result (2.3). □

Remark 1. If $0 < x < y$ and the function $f : [\ln x, \ln y] \rightarrow \mathbb{R}$ satisfies the condition (2.1) on the interval $[\ln x, \ln y]$, then we have the inequalities

$$\begin{aligned} \frac{m}{\alpha^2} (A(x^\alpha, y^\alpha) - L(x^\alpha, y^\alpha)) \\ \leq A(f(\ln x), f(\ln y)) - \frac{1}{\ln y - \ln x} \int_{\ln x}^{\ln y} f(t) dt \\ \leq \frac{M}{\alpha^2} (A(x^\alpha, y^\alpha) - L(x^\alpha, y^\alpha)) \quad (2.9) \end{aligned}$$

and

$$\begin{aligned} \frac{m}{\alpha^2} (L(x^\alpha, y^\alpha) - G(x^\alpha, y^\alpha)) \\ \leq \frac{1}{\ln y - \ln x} \int_{\ln x}^{\ln y} f(t) dt - f(\ln G(x, y)) \\ \leq \frac{M}{\alpha^2} (L(x^\alpha, y^\alpha) - G(x^\alpha, y^\alpha)), \quad (2.10) \end{aligned}$$

where $A(p, q) := \frac{p+q}{2}$ is the arithmetic mean, $G(p, q) := \sqrt{pq}$ is the geometric mean and $L(p, q) := \frac{p-q}{\ln p - \ln q}$ is the logarithmic mean.

We need the following result that is of interest itself. It provides lower and upper bounds for the Jensen's difference

$$\sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right)$$

in the case of twice differentiable functions whose second derivatives are bounded by exponentials as in (2.1).

Lemma 1. Let $f : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable function with the property that there exists the constants $\alpha, m, M \in \mathbb{R}$ with $\alpha \neq 0$, $m < M$ and such that (2.1) is valid.

Then for any $x_i \in [a, b]$ and $p_i \geq 0$ with $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n p_i = 1$ we have the inequalities

$$\begin{aligned} \frac{m}{\alpha^2} \left(\sum_{i=1}^n p_i e^{\alpha x_i} - e^{\alpha(\sum_{i=1}^n p_i x_i)} \right) \\ \leq \sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right) \\ \leq \frac{M}{\alpha^2} \left(\sum_{i=1}^n p_i e^{\alpha x_i} - e^{\alpha(\sum_{i=1}^n p_i x_i)} \right). \quad (2.11) \end{aligned}$$

Proof. Since the auxiliary function $g_{m,\alpha} : [a, b] \rightarrow \mathbb{R}$ given by $g_{m,\alpha}(t) := f(t) - \frac{m}{\alpha^2}e^{\alpha t}$ is convex, then by Jensen's inequality we have

$$\begin{aligned} 0 &\leq \sum_{i=1}^n p_i g_{m,\alpha}(x_i) - g_{m,\alpha}\left(\sum_{i=1}^n p_i x_i\right) \\ &= \sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right) \\ &\quad - \frac{m}{\alpha^2} \left(\sum_{i=1}^n p_i e^{\alpha x_i} - e^{\alpha(\sum_{i=1}^n p_i x_i)} \right) \end{aligned}$$

which produces the first inequality.

The second inequality follows in a similar way by employing the auxiliary function $g_{M,\alpha} : [a, b] \rightarrow \mathbb{R}$ given by $g_{M,\alpha}(t) := \frac{M}{\alpha^2}e^{\alpha t} - f(t)$. $\square$

Remark 2. If $0 < x < y$ and the function $f : [\ln x, \ln y] \rightarrow \mathbb{R}$ satisfies the condition (2.1) on the interval $[\ln x, \ln y]$, then we have the inequalities

$$\begin{aligned} \frac{m}{\alpha^2} \left(\sum_{i=1}^n p_i y_i^\alpha - \prod_{i=1}^n y_i^{\alpha p_i} \right) &\leq \sum_{i=1}^n p_i f(\ln y_i) - f\left(\ln \prod_{i=1}^n y_i^{p_i}\right) \\ &\leq \frac{M}{\alpha^2} \left(\sum_{i=1}^n p_i y_i^\alpha - \prod_{i=1}^n y_i^{\alpha p_i} \right), \end{aligned} \quad (2.12)$$

where $0 < x \leq y_i \leq y$ for $i \in \{1, \dots, n\}$.

Utilising the Jensen's type inequality (2.11) we are able to provide some upper and lower bounds for the difference of the integral means

$$\frac{1}{b-a} \int_a^b f(x) dx - \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f\left(\sum_{i=1}^n p_i x_i\right) dx_1 \dots dx_n$$

where $p_i > 0$ with $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n p_i = 1$.

Theorem 2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a twice differentiable function with the property that there exists the constants $\alpha, m, M \in \mathbb{R}$ with $\alpha \neq 0, m < M$ and such that (2.1) is valid.

Then for any $p_i > 0$ with $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n p_i = 1$ we have the inequalities

$$\begin{aligned} & \frac{m}{\alpha^2} \left(\frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - \frac{1}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \right) \\ & \leq \frac{1}{b-a} \int_a^b f(x) dx - \frac{1}{(b-a)^n} \int_a^b \dots \int_a^b f \left(\sum_{i=1}^n p_i x_i \right) dx_1 \dots dx_n \\ & \leq \frac{M}{\alpha^2} \left(\frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - \frac{1}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \right). \quad (2.13) \end{aligned}$$

Proof. We integrate the inequality (2.11) on $[a, b]^n$ to get

$$\begin{aligned} & \frac{m}{\alpha^2} \left(\sum_{i=1}^n p_i \int_a^b \dots \int_a^b e^{\alpha x_i} dx_1 \dots dx_n - \int_a^b \dots \int_a^b e^{\alpha(\sum_{i=1}^n p_i x_i)} dx_1 \dots dx_n \right) \quad (2.14) \\ & \leq \sum_{i=1}^n p_i \int_a^b \dots \int_a^b f(x_i) dx_1 \dots dx_n - \int_a^b \dots \int_a^b f \left(\sum_{i=1}^n p_i x_i \right) dx_1 \dots dx_n \\ & \leq \frac{M}{\alpha^2} \left(\sum_{i=1}^n p_i \int_a^b \dots \int_a^b e^{\alpha x_i} dx_1 \dots dx_n - \int_a^b \dots \int_a^b e^{\alpha(\sum_{i=1}^n p_i x_i)} dx_1 \dots dx_n \right). \end{aligned}$$

Observe that

$$\begin{aligned} \int_a^b \dots \int_a^b e^{\alpha x_i} dx_1 \dots dx_n &= (b-a)^{n-1} \int_a^b e^{\alpha x_i} dx_i \\ &= (b-a)^n \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)}, \end{aligned}$$

$$\begin{aligned} \int_a^b \dots \int_a^b e^{\alpha(\sum_{i=1}^n p_i x_i)} dx_1 \dots dx_n &= \int_a^b \dots \int_a^b \prod_{i=1}^n e^{\alpha p_i x_i} dx_1 \dots dx_n \\ &= \prod_{i=1}^n \int_a^b e^{\alpha p_i x_i} dx_i = \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{\alpha p_i} \\ &= \frac{(b-a)^n}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \end{aligned}$$

and

$$\int_a^b \dots \int_a^b f(x_i) dx_1 \dots dx_n = (b-a)^{n-1} \int_a^b f(x) dx.$$

From (2.14) we then get

$$\begin{aligned}
 & \frac{m}{\alpha^2} \left((b-a)^n \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - \frac{(b-a)^n}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \right) \\
 & \leq (b-a)^{n-1} \int_a^b f(x) dx - \int_a^b \dots \int_a^b f \left(\sum_{i=1}^n p_i x_i \right) dx_1 \dots dx_n \\
 & \leq \frac{M}{\alpha^2} \left((b-a)^n \frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - \frac{(b-a)^n}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \right),
 \end{aligned}$$

which by division with $(b-a)^n$ produces the desired result (2.13). $\square$

3. SOME APPLICATIONS

The above inequalities may be applied for various functions in Analysis for which simple upper and lower bounds for the function $\frac{f''(\cdot)}{e^{\alpha \cdot}}$ can be found.

Consider, for instance, the function $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ given by $f(t) = \frac{1}{t}$.

We have $f''(t) = \frac{2}{t^3}$ for $t \in [a, b]$ and

$$\frac{2}{b^3 e^b} \leq \frac{f''(t)}{e^t} \leq \frac{2}{a^3 e^a} \quad (3.1)$$

for any $t \in [a, b]$.

Utilising the inequality (2.2) we obtain

$$\begin{aligned}
 \frac{2}{b^3 e^b} (A(e^a, e^b) - L(e^a, e^b)) & \leq \frac{L(a, b) - H(a, b)}{L(a, b) H(a, b)} \\
 & \leq \frac{2}{a^3 e^a} (A(e^a, e^b) - L(e^a, e^b))
 \end{aligned} \quad (3.2)$$

and from (2.3)

$$\begin{aligned}
 \frac{2}{b^3 e^b} (L(e^a, e^b) - e^{A(a, b)}) & \leq \frac{A(a, b) - L(a, b)}{A(a, b) L(a, b)} \\
 & \leq \frac{2}{a^3 e^a} (L(e^a, e^b) - e^{A(a, b)}),
 \end{aligned} \quad (3.3)$$

where $H(a, b) := \frac{2}{\frac{1}{a} + \frac{1}{b}}$ is the harmonic mean.

Now, consider the function $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ given by $f(t) = e^{\beta t}$ with $\beta > \alpha$. Then we have

$$\beta^2 e^{(\beta-\alpha)a} \leq \frac{f''(t)}{e^{\alpha t}} \leq \beta^2 e^{(\beta-\alpha)b} \quad (3.4)$$

for any $t \in [a, b]$.

If we apply the inequality (2.11) for the function $f(t) = e^{\beta t}$, $t \in [a, b]$, then we have the inequalities

$$\begin{aligned} \frac{\beta^2}{\alpha^2} e^{(\beta-\alpha)a} \left(\sum_{i=1}^n p_i e^{\alpha x_i} - e^{\alpha(\sum_{i=1}^n p_i x_i)} \right) \\ \leq \sum_{i=1}^n p_i e^{\beta x_i} - e^{\beta(\sum_{i=1}^n p_i x_i)} \\ \leq \frac{\beta^2}{\alpha^2} e^{(\beta-\alpha)b} \left(\sum_{i=1}^n p_i e^{\alpha x_i} - e^{\alpha(\sum_{i=1}^n p_i x_i)} \right) \end{aligned} \quad (3.5)$$

for any $x_i \in [a, b]$ and $p_i \geq 0$ with $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n p_i = 1$.

Now, assume that $0 < s \leq y_i \leq S < \infty$ for any $i \in \{1, \dots, n\}$. On choosing $x_i = \ln y_i$ for any $i \in \{1, \dots, n\}$, then we have $\ln s \leq x_i \leq \ln S$.

If we write the inequality (3.5) for these $x_i = \ln y_i$ for any $i \in \{1, \dots, n\}$ we get for $\beta > \alpha$

$$\begin{aligned} \frac{\beta^2}{\alpha^2} s^{(\beta-\alpha)} \left(\sum_{i=1}^n p_i y_i^\alpha - \prod_{i=1}^n y_i^{\alpha p_i} \right) &\leq \sum_{i=1}^n p_i y_i^\beta - \prod_{i=1}^n y_i^{\beta p_i} \\ &\leq \frac{\beta^2}{\alpha^2} S^{(\beta-\alpha)} \left(\sum_{i=1}^n p_i y_i^\alpha - \prod_{i=1}^n y_i^{\alpha p_i} \right) \end{aligned} \quad (3.6)$$

provided that $0 < s \leq y_i \leq S < \infty$, $p_i \geq 0$ for any $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n p_i = 1$.

If in this inequality we take $\beta = 1$ and $\alpha = -1$ then we get

$$s^2 \left(\sum_{i=1}^n \frac{p_i}{y_i} - \frac{1}{\prod_{i=1}^n y_i^{p_i}} \right) \leq \sum_{i=1}^n p_i y_i - \prod_{i=1}^n y_i^{p_i} \leq S^2 \left(\sum_{i=1}^n \frac{p_i}{y_i} - \frac{1}{\prod_{i=1}^n y_i^{p_i}} \right). \quad (3.7)$$

Finally, on applying the inequality (2.13) for the exponential function $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ given by $f(t) = e^{\beta t}$ with $\beta > \alpha$, we obtain

$$\begin{aligned} \frac{\beta^2}{\alpha^2} e^{(\beta-\alpha)a} \left(\frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - \frac{1}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \right) \\ \leq \frac{e^{\beta b} - e^{\alpha a}}{\beta(b-a)} - \frac{1}{\beta^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\beta p_i b} - e^{\beta p_i a}}{b-a} \\ \leq \frac{\beta^2}{\alpha^2} e^{(\beta-\alpha)b} \left(\frac{e^{\alpha b} - e^{\alpha a}}{\alpha(b-a)} - \frac{1}{\alpha^n \prod_{i=1}^n p_i} \prod_{i=1}^n \frac{e^{\alpha p_i b} - e^{\alpha p_i a}}{b-a} \right) \end{aligned} \quad (3.8)$$

for any $p_i > 0$ with $i \in \{1, \dots, n\}$ and $\sum_{i=1}^n p_i = 1$.

REFERENCES

- [1] Allasia, G., Giordano, C., & Pečarić, J. (1999). Hadamard-type inequalities for $(2r)$ -convex functions with applications. *Atti Acad. Sci. Torino-Cl. Sc. Fis.*, 133, 1-14.
- [2] Alzer, H. (1989a). A note on Hadamard's inequalities. *C. R. Math. Rep. Acad. Sci. Canada*, 11, 255-258.
- [3] Alzer, H. (1989b). On an integral inequality. *Math. Rev. Anal. Numer. Theor. Approx.*, 18, 101-103.
- [4] Azpeitia, A. G. (1994). Convex functions and the Hadamard inequality. *Rev.-Colombiana-Mat.*, 28(1), 7-12.
- [5] Barbu, D., Dragomir, S. S., & Buşe, C. (1993). A probabilistic argument for the convergence of some sequences associated to Hadamard's inequality. *Studia Univ. Babeş-Bolyai. Math.*, 38(1), 29-33.
- [6] Buşe, C., Dragomir, S. S., & Barbu, D. (1996). The convergence of Some Sequences Connected to Hadamard's inequality. *Demonstratio Math.*, 29(1), 53-59.
- [7] Dragomir, S. S. (1991). A mapping in connection to Hadamard's inequalities. *An. Öster. Akad. Wiss. Math.-Natur. (Wien)*, 128, 17-20.
- [8] Dragomir, S. S. (1992a). On Hadamard's inequalities for convex functions. *Mat. Balkanica*, 6, 215-222.
- [9] Dragomir, S. S. (1992b). Some integral inequalities for differentiable convex functions. *Contributions, Macedonian Acad. of Sci. and Arts*, 13(1), 13-17.
- [10] Dragomir, S. S. (1992c). Two mappings in connection to Hadamard's inequalities. *J. Math. Anal. Appl.*, 167, 49-56.
- [11] Dragomir, S. S. (1993). A Refinement of Hadamard's inequality for isotonic linear functionals. *Tamkang J. of Math. (Taiwan)*, 24, 101-106.
- [12] Dragomir, S. S. (1994). Some remarks on Hadamard's inequalities for convex functions. *Extracta Math.*, 9(2), 88-94.
- [13] Dragomir, S. S. (2000a). On Hadamard's inequality for the convex mappings defined on a ball in the space and applications. *Math. Ineq. & Appl.*, 3(2), 177-187.
- [14] Dragomir, S. S. (2000b). On Hadamard's inequality on a disk. *Journal of Ineq. in Pure & Appl. Math.*, 1(1), Article 2, 1-11.
- [15] Dragomir, S. S., & Agarwal, R. P. (1998). Two new mappings associated with Hadamard's inequalities for convex functions. *Appl. Math. Lett.*, 11(3), 33-38.
- [16] Dragomir, S. S., & Buşe, C. (1995). Refinements of Hadamard's inequality for multiple integrals. *Utilitas Math. (Canada)*, 47, 193-195.
- [17] Dragomir, S. S., Cho, Y. J., & Kim, S. S. (2000). Inequalities of Hadamard's type for lipschitzian mappings and their applications. *J. of Math. Anal. Appl.*, 245(2), 489-501.
- [18] Dragomir, S. S., & Fitzpatrick, S. (1998). The Hadamard's inequality for s -convex functions in the first sense. *Demonstratio Math.*, 31(3), 633-642.
- [19] Dragomir, S. S., & Fitzpatrick, S. (1999). The Hadamard's inequality for s -convex functions in the second sense. *Demonstratio Math.*, 32(4), 687-696.

- [20] Dragomir, S. S., & Gomm, I. (2011). Bounds for two mappings associated to the Hermite-Hadamard inequality. *Aust. J. Math. Anal. Appl.*, 8, Article 5, 1-9.
- [21] Dragomir, S. S., & Gomm, I. (2012). Some new bounds for two mappings related to the Hermite-Hadamard inequality for convex functions. *Num. Alg. Cont. & Opt.*, 2(2), 271-278.
- [22] Dragomir, S. S., & Ionescu, N. M. (1990). On some inequalities for convex-dominated functions. *Anal. Num. Theor. Approx.*, 19, 21-28.
- [23] Dragomir, S. S., Milošević, D. S., & Sándor, J. (1993). On some refinements of Hadamard's inequalities and applications. *Univ. Belgrad, Publ. Elek. Fak. Sci. Math.*, 4, 21-24.
- [24] Dragomir, S. S., & Mond, B. (1997). On Hadamard's inequality for a class of functions of Godunova and Levin. *Indian J. Math.*, 39(1), 1-9.
- [25] Dragomir, S. S., & Pearce, C. E. M. (1998). Quasi-convex functions and Hadamard's inequality. *Bull. Austral. Math. Soc.*, 57, 377-385.
- [26] Dragomir, S. S., & Pearce, C. E. M. (2000). *Selected topics on Hermite-Hadamard inequalities and applications*. RGMIA Monographs, Retrieved from http://rgmia.org/monographs/hermite_hadamard.html
- [27] Dragomir, S. S., Pearce, C. E. M., & Pečarić, J. (1995a). On Jessen's and related inequalities for isotonic sublinear functionals. *Acta Math. Sci. (Szeged)*, 61, 373-382.
- [28] Dragomir, S. S., Pečarić, J., & Persson, L. E. (1995b). Some inequalities of Hadamard type. *Soochow J. of Math. (Taiwan)*, 21, 335-341.
- [29] Dragomir, S. S., Pečarić, J. & Sándor, J. (1990). A note on the Jensen-Hadamard inequality. *Anal. Num. Theor. Approx.*, 19, 21-28.
- [30] Dragomir, S. S., & Toader, G. H. (1993). Some inequalities for m -convex functions. *Studia Univ. Babeş-Bolyai, Math.*, 38(1), 21-28.
- [31] Fink, A. M. (1994). Toward a theory of best possible inequalities. *Nieuw Archief von Wiskunde*, 12, 19-29.
- [32] Fink, A. M. (1995). Two inequalities. *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat.*, 6, 48-49.
- [33] Fink, A. M. (1998). A best possible Hadamard inequality. *Math. Ineq. & Appl.*, 2, 223-230.
- [34] Gavrea, B. (2000). On Hadamard's inequality for the convex mappings defined on a convex domain in the space. *Journal of Ineq. in Pure & Appl. Math.*, 1(1), Article 9, 1-6.
- [35] Gill, P. M., Pearce, C. E. M., & Pečarić, J. (1997). Hadamard's inequality for r -convex functions. *Journal of Math. Anal. and Appl.*, 215, 461-470.
- [36] Hardy, G. H., Littlewood, E. J., & Pólya, G. (1952). *Inequalities* (2nd ed.). Cambridge University Press.
- [37] Lee, K.-C., & Tseng, K.-L. (2000). On a weighted generalisation of Hadamard's inequality for G -convex functions. *Tamsui Oxford Journal of Math. Sci.*, 166(1), 91-104.
- [38] Lupaş, A. (1976). A generalisation of Hadamard's inequality for convex functions. *Univ. Beograd. Publ. Elek. Fak. Ser. Mat. Fiz.*, (544-576), 115-121.
- [39] Lupaş, A. (1997). The Jensen-Hadamard inequality for convex functions of higher order. *Octagon Math. Mag.*, 5(2), 8-9.
- [40] Maksimović, D. M. (1979). A short proof of generalized Hadamard's

- inequalities. *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, (634-677), 126-128.
- [41] Mitrinović, D. S., & Lacković, I. (1985). Hermite and convexity. *Aequat. Math.*, 28, 229-232.
 - [42] Mitrinović, D. S., Pečarić, J., & Fink, A. M. (1993). *Classical and new inequalities in analysis*. Kluwer Academic Publishers, Dordrecht/Boston/London.
 - [43] Neuman, E. (1986). Inequalities involving generalised symmetric means. *J. Math. Anal. Appl.*, 120, 315-320.
 - [44] Neuman, E., & Pečarić, J. (1989). Inequalities involving multivariate convex functions. *J. Math. Anal. Appl.*, 137, 514-549.
 - [45] Neuman, E. (1990). Inequalities involving multivariate convex functions II. *Proc. Amer. Math. Soc.*, 190, 965-974.
 - [46] Niculescu, C. P. (2000a). A note on the dual Hermite-Hadamard inequality. *The Math. Gazette*, (Romania), July, 1-3.
 - [47] Niculescu, C. P. (2000b). Convexity according to the geometric mean. *Math. Ineq. & Appl.*, 3(2), 155-167.
 - [48] Pearce, C. E. M., Pečarić, J., & Šimić, V. (1998). Stolarsky means and Hadamard's inequality. *J. Math. Anal. Appl.*, 220, 99-109.
 - [49] Pearce, C. E. M., & Rubinov, A. M. (1999). P -functions, quasi-convex functions and Hadamard-type inequalities. *J. Math. Anal. Appl.*, 240(1), 92-104.
 - [50] Pečarić, J. (1992). Remarks on two interpolations of Hadamard's inequalities. *Contributions, Macedonian Acad. of Sci. and Arts, Sect. of Math. and Technical Sciences, (Scopje)*, 13, 9-12.
 - [51] Pečarić, J., & Dragomir, S. S. (1991). A generalization of Hadamard's integral inequality for isotonic linear functionals. *Rudovi Mat. (Sarajevo)*, 7, 103-107.
 - [52] Pečarić, J., Proschan, F., & Tong, Y. L. (1992). *Convex functions, partial orderings and statistical applications*. Academic Press, Inc..
 - [53] Sándor, J. (1990a). A note on the Jensen-Hadamard inequality. *Anal. Numer. Theor. Approx.*, 19(1), 29-34.
 - [54] Sándor, J. (1990b). An application of the Jensen-Hadamard inequality. *Nieuw-Arch.-Wisk.*, 8(1), 63-66.
 - [55] Sándor, J. (1991). On the Jensen-Hadamard inequality. *Studia Univ. Babes-Bolyai, Math.*, 36(1), 9-15.
 - [56] Vasić, P. M., Lacković, I. B., & Maksimović, D. M. (1980). Note on convex functions IV: on Hadamard's inequality for weighted arithmetic means. *Univ. Beograd Publ. Elek. Fak., Ser. Mat. Fiz.*, (678-715), 199-205.
 - [57] Yang, G. S., & Hong, M. C. (1997). A note on Hadamard's inequality. *Tamkang J. Math.*, 28(1), 33-37.
 - [58] Yang, G. S., & Tseng, K. L. (1999). On certain integral inequalities related to Hermite-Hadamard inequalities. *J. Math. Anal. Appl.*, 239, 180-187.